

Signal Processing First

Lecture 6 Fourier Series Coefficients

9/8/2003

© 2003, JH McClellan & RW Schafer

1

READING ASSIGNMENTS

- This Lecture:
 - **Fourier Series in Ch 3, Sects 3-4, 3-5 & 3-6**
 - Replaces pp. 62-66 in Ch 3 in DSP First
 - Notation: a_k for Fourier Series
- Other Reading:
 - Next Lecture: More Fourier Series

9/8/2003

© 2003, JH McClellan & RW Schafer

3

LECTURE OBJECTIVES

- Work with the Fourier Series Integral

$$a_k = \frac{1}{T_0} \int_0^{T_0} x(t) e^{-j(2\pi k/T_0)t} dt$$

- **ANALYSIS** via Fourier Series
 - For **PERIODIC** signals: $x(t+T_0) = x(t)$
 - Later: spectrum from the Fourier Series

9/8/2003

© 2003, JH McClellan & RW Schafer

4

HISTORY

- Jean Baptiste Joseph Fourier
 - 1807 thesis (memoir)
 - On the Propagation of Heat in Solid Bodies
 - Heat !
 - Napoleonic era
- <http://www-groups.dcs.st-and.ac.uk/~history/Mathematicians/Fourier.html>

9/8/2003

© 2003, JH McClellan & RW Schafer

5




Joseph Fourier

lived from 1768 to 1830

Fourier studied the mathematical theory of heat conduction. He established the partial differential equation governing heat diffusion and solved it by using infinite series of trigonometric functions.

SPECTRUM DIAGRAM

- Recall Complex Amplitude vs. Freq



Harmonic Signal

$$x(t) = \sum_{k=-\infty}^{\infty} a_k e^{j2\pi k f_0 t}$$

PERIOD/FREQUENCY of COMPLEX EXPONENTIAL:

$$2\pi(f_0) = \omega_0 = \frac{2\pi}{T_0} \quad \text{or} \quad T_0 = \frac{1}{f_0}$$

Fourier Series Synthesis

$$x(t) = \sum_{k=-\infty}^{\infty} a_k e^{j2\pi k f_0 t}$$


$$a_k = \frac{1}{2} X_k = \frac{1}{2} A_k e^{j\phi_k}$$

$$x(t) = A_0 + \sum_{k=1}^N A_k \cos(2\pi k f_0 t + \phi_k)$$

$$X_k = A_k e^{j\phi_k}$$

COMPLEX AMPLITUDE

Harmonic Signal (3 Freqs)



SYNTHESIS vs. ANALYSIS

- SYNTHESIS
 - Easy
 - Given (ω_k, A_k, ϕ_k) create $x(t)$
- ANALYSIS
 - Hard
 - Given $x(t)$, extract (ω_k, A_k, ϕ_k)
 - How many?
 - Need algorithm for computer
- Synthesis can be HARD
 - Synthesize Speech so that it sounds good

9/8/2003

© 2003, JH McClellan & RW Schafer

11

STRATEGY: $x(t) \rightarrow a_k$

- ANALYSIS
 - Get representation from the signal
 - Works for **PERIODIC** Signals
- Fourier Series
 - Answer is: an INTEGRAL over one period

$$a_k = \frac{1}{T_0} \int_0^{T_0} x(t) e^{-j\omega_0 k t} dt$$

INTEGRAL Property of exp(j)

- INTEGRATE over ONE PERIOD

$$\int_0^{T_0} e^{-j(2\pi/T_0)mt} dt = \frac{T_0}{-j2\pi m} e^{-j(2\pi/T_0)mt} \Big|_0^{T_0}$$

$$= \frac{T_0}{-j2\pi m} (e^{-j2\pi m} - 1)$$

$$\int_0^{T_0} e^{-j(2\pi/T_0)mt} dt = 0$$

$m \neq 0$

$$\omega_0 = \frac{2\pi}{T_0}$$

9/8/2003

© 2003, JH McClellan & RW Schafer

12

9/8/2003

© 2003, JH McClellan & RW Schafer

13

ORTHOGONALITY of exp(j)

- PRODUCT of exp(+j) and exp(-j)

$$\frac{1}{T_0} \int_0^{T_0} e^{j(2\pi/T_0)\ell t} e^{-j(2\pi/T_0)kt} dt = \begin{cases} 0 & k \neq \ell \\ 1 & k = \ell \end{cases}$$

$$\frac{1}{T_0} \int_0^{T_0} e^{j(2\pi/T_0)(\ell-k)t} dt$$

9/8/2003

© 2003, JH McClellan & RW Schaffer

14

Isolate One FS Coefficient

$$x(t) = \sum_{k=-\infty}^{\infty} a_k e^{j(2\pi/T_0)kt}$$

$$\frac{1}{T_0} \int_0^{T_0} x(t) e^{-j(2\pi/T_0)\ell t} dt = \frac{1}{T_0} \int_0^{T_0} \left(\sum_{k=-\infty}^{\infty} a_k e^{j(2\pi/T_0)kt} \right) e^{-j(2\pi/T_0)\ell t} dt$$

$$\frac{1}{T_0} \int_0^{T_0} x(t) e^{-j(2\pi/T_0)\ell t} dt = \sum_{k=-\infty}^{\infty} a_k \left(\frac{1}{T_0} \int_0^{T_0} e^{j(2\pi/T_0)(k-\ell)t} dt \right) = a_\ell$$

Integral is zero except for $k = \ell$

$$\Rightarrow a_k = \frac{1}{T_0} \int_0^{T_0} x(t) e^{-j(2\pi/T_0)kt} dt$$

9/8/2003


© 2003, JH McClellan & RW Schaffer

15

SQUARE WAVE EXAMPLE

$$x(t) = \begin{cases} 1 & 0 \leq t < \frac{1}{2} T_0 \\ 0 & \frac{1}{2} T_0 \leq t < T_0 \end{cases}$$

for $T_0 = 0.04$ sec.



9/8/2003

© 2003, JH McClellan & RW Schaffer

16

FS for a SQUARE WAVE {a_k}

$$a_k = \frac{1}{T_0} \int_0^{T_0} x(t) e^{-j(2\pi/T_0)kt} dt \quad (k \neq 0)$$

$$a_k = \frac{1}{.04} \int_0^{.02} 1 e^{-j(2\pi/.04)kt} dt = \frac{1}{.04(-j2\pi k/.04)} e^{-j(2\pi/.04)kt} \Big|_0^{.02}$$

$$= \frac{1}{(-j2\pi k)} (e^{-j(\pi)k} - 1) = \frac{1 - (-1)^k}{j2\pi k}$$

9/8/2003

© 2003, JH McClellan & RW Schaffer

17

DC Coefficient: a_0

$$a_k = \frac{1}{T_0} \int_0^{T_0} x(t) e^{-j(2\pi/T_0)kt} dt \quad (k = 0)$$

$$a_0 = \frac{1}{T_0} \int_0^{T_0} x(t) dt = \frac{1}{T_0} (\text{Area})$$

$$a_0 = \frac{1}{.04} \int_0^{.02} 1 dt = \frac{1}{.04} (.02 - 0) = \frac{1}{2}$$

9/8/2003

© 2003, JH McClellan & RW Schaefer

18

Fourier Coefficients a_k

- a_k is a function of k
 - Complex Amplitude for k -th Harmonic
 - This one doesn't depend on the period, T_0

$$a_k = \frac{1 - (-1)^k}{j2\pi k} = \begin{cases} \frac{1}{j\pi k} & k = \pm 1, \pm 3, \dots \\ 0 & k = \pm 2, \pm 4, \dots \\ \frac{1}{2} & k = 0 \end{cases}$$

9/8/2003


© 2003, JH McClellan & RW Schaefer

19

Spectrum from Fourier Series

$$\omega_0 = 2\pi / (0.04) = 2\pi(25)$$

$$a_k = \begin{cases} \frac{-j}{\pi k} & k = \pm 1, \pm 3, \dots \\ 0 & k = \pm 2, \pm 4, \dots \\ \frac{1}{2} & k = 0 \end{cases}$$



9/8/2003

© 2003, JH McClellan & RW Schaefer

20

Fourier Series Integral

- HOW do you determine a_k from $x(t)$?

$$a_k = \frac{1}{T_0} \int_0^{T_0} x(t) e^{-j(2\pi/T_0)kt} dt$$

Fundamental Frequency $f_0 = 1/T_0$

$a_{-k} = a_k^*$ when $x(t)$ is real

$$a_0 = \frac{1}{T_0} \int_0^{T_0} x(t) dt \quad (\text{DC component})$$

9/8/2003

© 2003, JH McClellan & RW Schaefer

21